## AUTHOR INDEX

Aiuti, F. see Poli, G. Acero, R. see Poli, G. Acchain, D. see Santoro, F. Aizawa, M. see Ishikura, H. Akatsuka, T. see Kubota, T.

AKBAR, A.N., GIARDINA, P.J., HILGARTNER, M.W. & GRAY, R.W. Immunoglobulin abnormalities in thalassaemia major. I. A transfusion-related increase in circulating cytoplasmic immunoglobulin-positive cells, 397

VAN DEN AKKER, T.W. see RADL, J.

VAN OUD ALBLAS, A.B., JANSSENS, A.R., LEIJH, P.C.J. & van Furth, R. Functions of granulocytes and monocytes in primary biliary and alcoholic cirrhosis, 724

AL-SAKKAF, L., POZZILLI, P., SENSI, M., IRVING, W.L. & BOTTAZZO, G.F. A cell surface monoclonal antibody (H366) helps to discriminate human cytotoxic from suppressor T cells, 594

ANTONACI, S., JIRILLO, E., KIYONO, H., WILLIAMSON, S.I., MICHALEK, S.M. & MCGHEE. Relationship between immune system and gram-negative bacteria. III. Functional analysis of human peripheral blood lymphocyte subpopulations separated by cytoadherence with Salmonella minnesota Rb, 248

VAN ÁSPEREN, P.P., GLEESON, M., KEMP, A.S., CRIPPS, A.W., GERAGHTY, S.B., MELLIS, C.M. & CLANCY, R.L. The relationship between atopy and salivary IgA deficiency in infancy, 753

ATKINSON, J.P. see SEYA, T.

AUWERX, J., BOOGAERTS, M., CEUPPENS, J.L. & DEMEDTS. Defective host defence mechanisms in a family with hypocalciuric hypercalcaemia and coexisting interstitial lung disease, 57

BACON, P.A. see SALMON, M. BENNER, R. see RADL. J.

BERG. P.A. see FALCIONI, F. BERSANI, L. see POLI, G.

BETTERLE, C., CARETTO, A., ZEVIANI, M., PEDINI, B. & SALVIATI, C. Demonstration and characterization of anti-human mitochondria autoantibodies in idiopathic hypoparathyroidism and in other conditions. 353

BIBERFELD, P. see ROBERT, K.-H. BLANN, A.D. see Salmon, M.

BLITSTEIN-WILLINGER, E. Normalization of defective interleukin 1 and interleukin 2 production in patients with pemphigus vulgaris following chrysotherapy, 705

BOCCHIERI, M.H., SMITH JR, J.B., SMITH, J.B., J. STARUCH, M.J. & WOOD, D.D. Functional consequences of perceived interleukin deficiencies? Analysis employing NZB×C58 recombinant inbred mice, 622

BOFILL, M. see PARKHOUSE, R.M.E. BOOGAERTS, M. see AUWERX, J. BORASCHI, D. see TAGLIAUBE, A.

BOTTAZZI, B. see POLI, G.

BOTTAZZO, G.F. see AL-SAKKAF, L.

BOURASSA, D., FORGET, A., PELLETIER, M., SKAMENE, E. & TURCOTTE, R. Cellular immune response to

Mycobacterium bovis (BCG) in genetically-susceptible and resistant congenic mouse strains, 31

BOYLE, C.C. see LOIZOU, S.

Brennand, D.M. see Maddison, P.J.

BRESNIHAN, B. see HANLY, J.G.
BROWN, V.A., SMITH, S.K., DEWAR, A.E., STOCKDILL,
G. & MADDY, A.H. Surface glycoproteins ass
markers of the cellular status of B chronic
lymphocytic leukaemia lymphocytes, 95

BRULLEY-ROSSET, M., DARDENNE, M. & SCHUURS, A. Functional and quantitative changes of immune cells of ageing NZB mice treated with nandrolone decanoate. I. Effect on survival and autoantibody development, 630

CAFFARENA, A. see TAGLIAUBE, A.
CALVALIERI, S. see TAGLIAUBE, A.
VAN CAMP, B. see DE WAELE, M.
CAPRON, A. see SANTORO, F.
CARBONARI, M. see POLI, G.
CARETTO, A. see BETTERLE, C.
CARSON DICK, W. see GEORGOPOULOS, K.
CAVENDER, D. see YU, C.-L.

CAZZOLA, G. see TAGLIAUBE, A. CESBRON, J.Y. see SANTORO, F.

CEUPPENS, J.L. see AUWERX, J. CHRISTIANSEN, F.T. see UKO, G.

CLANCY, R.L. see VAN ASPEREN, P.P. CLAUDE, P. see VUITTON, D.A.

COHEN, P.L. see EISENBERG, R.A.
COLLINGS, L.A., WATERS, M.F.R. & POULTER, L.W.
The involvement of dendritic cells in the cutaneous lesions associated with tuberculoid and lepromatous leprosy, 458

COOPER, D.A. see ZAUNDERS, J.J. CRIPPS, A.W. see VAN ASPEREN, P.P.

CUKIER, R. & TRON, F. Monoclonal anti-DNA antibodies: an approach to studying SLE nephritis, 143

D'Amoro, J.D. see Wursch, A.M. Das, P.K. see Roozemon, R.C.

DAWKINS, R.L. see UKO, G.

DE AZIPURUA, H.J. see Dow, C.A.

DE CARVALHO, L.C.P. see KRETTLI, A.U.

DE GAST see WURSCH, A.M. DE SOUZA, J.B. see PLAYFAIR, J.H.

DE WAELE, M., NAESSENS, A., FOULON, W. & VAN CAMP, B. Activated T-cells with suppressor/cytotoxic phenotype in acute *Toxoplasma gondii* infection, 256

DE WIT, Y.L. see KLASTER, P.R. DEGEBRODT, A. see HOLY, P.G.

DELACROIX, D.L. see VUITTON, D.A.

DEMEDTS, M. see AUWERX, J. DENMAN, A.M. see PELTON, B.K.

DERNICK, R. see UHLIG, H. DEWAR, A.E. see BROWN, V.A.

DIESFELD, H.J. see RUPPEL, A.

DILLY, S.A. & SLOANE, J.P. An immunohistological study of human nephrotic graft-versus-host disease. 545

## AUTHOR INDEX

Aiuti, F. see Poli, G. Acero, R. see Poli, G. Acchain, D. see Santoro, F. Aizawa, M. see Ishikura, H. Akatsuka, T. see Kubota, T.

AKBAR, A.N., GIARDINA, P.J., HILGARTNER, M.W. & GRAY, R.W. Immunoglobulin abnormalities in thalassaemia major. I. A transfusion-related increase in circulating cytoplasmic immunoglobulin-positive cells, 397

VAN DEN AKKER, T.W. see RADL, J.

VAN OUD ALBLAS, A.B., JANSSENS, A.R., LEIJH, P.C.J. & van Furth, R. Functions of granulocytes and monocytes in primary biliary and alcoholic cirrhosis, 724

AL-SAKKAF, L., POZZILLI, P., SENSI, M., IRVING, W.L. & BOTTAZZO, G.F. A cell surface monoclonal antibody (H366) helps to discriminate human cytotoxic from suppressor T cells, 594

ANTONACI, S., JIRILLO, E., KIYONO, H., WILLIAMSON, S.I., MICHALEK, S.M. & MCGHEE. Relationship between immune system and gram-negative bacteria. III. Functional analysis of human peripheral blood lymphocyte subpopulations separated by cytoadherence with Salmonella minnesota Rb, 248

VAN ÁSPEREN, P.P., GLEESON, M., KEMP, A.S., CRIPPS, A.W., GERAGHTY, S.B., MELLIS, C.M. & CLANCY, R.L. The relationship between atopy and salivary IgA deficiency in infancy, 753

ATKINSON, J.P. see SEYA, T.

AUWERX, J., BOOGAERTS, M., CEUPPENS, J.L. & DEMEDTS. Defective host defence mechanisms in a family with hypocalciuric hypercalcaemia and coexisting interstitial lung disease, 57

BACON, P.A. see SALMON, M. BENNER, R. see RADL. J.

BERG. P.A. see FALCIONI, F. BERSANI, L. see POLI, G.

BETTERLE, C., CARETTO, A., ZEVIANI, M., PEDINI, B. & SALVIATI, C. Demonstration and characterization of anti-human mitochondria autoantibodies in idiopathic hypoparathyroidism and in other conditions. 353

BIBERFELD, P. see ROBERT, K.-H. BLANN, A.D. see Salmon, M.

BLITSTEIN-WILLINGER, E. Normalization of defective interleukin 1 and interleukin 2 production in patients with pemphigus vulgaris following chrysotherapy, 705

BOCCHIERI, M.H., SMITH JR, J.B., SMITH, J.B., J. STARUCH, M.J. & WOOD, D.D. Functional consequences of perceived interleukin deficiencies? Analysis employing NZB×C58 recombinant inbred mice, 622

BOFILL, M. see PARKHOUSE, R.M.E. BOOGAERTS, M. see AUWERX, J. BORASCHI, D. see TAGLIAUBE, A.

BOTTAZZI, B. see POLI, G.

BOTTAZZO, G.F. see AL-SAKKAF, L.

BOURASSA, D., FORGET, A., PELLETIER, M., SKAMENE, E. & TURCOTTE, R. Cellular immune response to

Mycobacterium bovis (BCG) in genetically-susceptible and resistant congenic mouse strains, 31

BOYLE, C.C. see LOIZOU, S.

Brennand, D.M. see Maddison, P.J.

BRESNIHAN, B. see HANLY, J.G.
BROWN, V.A., SMITH, S.K., DEWAR, A.E., STOCKDILL,
G. & MADDY, A.H. Surface glycoproteins ass
markers of the cellular status of B chronic
lymphocytic leukaemia lymphocytes, 95

BRULLEY-ROSSET, M., DARDENNE, M. & SCHUURS, A. Functional and quantitative changes of immune cells of ageing NZB mice treated with nandrolone decanoate. I. Effect on survival and autoantibody development, 630

CAFFARENA, A. see TAGLIAUBE, A.
CALVALIERI, S. see TAGLIAUBE, A.
VAN CAMP, B. see DE WAELE, M.
CAPRON, A. see SANTORO, F.
CARBONARI, M. see POLI, G.
CARETTO, A. see BETTERLE, C.
CARSON DICK, W. see GEORGOPOULOS, K.
CAVENDER, D. see YU, C.-L.

CAZZOLA, G. see TAGLIAUBE, A. CESBRON, J.Y. see SANTORO, F.

CEUPPENS, J.L. see AUWERX, J. CHRISTIANSEN, F.T. see UKO, G.

CLANCY, R.L. see VAN ASPEREN, P.P. CLAUDE, P. see VUITTON, D.A.

COHEN, P.L. see EISENBERG, R.A.
COLLINGS, L.A., WATERS, M.F.R. & POULTER, L.W.
The involvement of dendritic cells in the cutaneous lesions associated with tuberculoid and lepromatous leprosy, 458

COOPER, D.A. see ZAUNDERS, J.J. CRIPPS, A.W. see VAN ASPEREN, P.P.

CUKIER, R. & TRON, F. Monoclonal anti-DNA antibodies: an approach to studying SLE nephritis, 143

D'Amoro, J.D. see Wursch, A.M. Das, P.K. see Roozemon, R.C.

DAWKINS, R.L. see UKO, G.

DE AZIPURUA, H.J. see Dow, C.A.

DE CARVALHO, L.C.P. see KRETTLI, A.U.

DE GAST see WURSCH, A.M. DE SOUZA, J.B. see PLAYFAIR, J.H.

DE WAELE, M., NAESSENS, A., FOULON, W. & VAN CAMP, B. Activated T-cells with suppressor/cytotoxic phenotype in acute *Toxoplasma gondii* infection, 256

DE WIT, Y.L. see KLASTER, P.R. DEGEBRODT, A. see HOLY, P.G.

DELACROIX, D.L. see VUITTON, D.A.

DEMEDTS, M. see AUWERX, J. DENMAN, A.M. see PELTON, B.K.

DERNICK, R. see UHLIG, H. DEWAR, A.E. see BROWN, V.A.

DIESFELD, H.J. see RUPPEL, A.

DILLY, S.A. & SLOANE, J.P. An immunohistological study of human nephrotic graft-versus-host disease. 545 Dow, C.A., De AZIPURUA, H.J., PEDERSEN, J.S., UNGAR, B. & TOH, B.H. 65-75 kD protein identified by immunoblotting as the presumptive gastric microsomal autoantigen in pernicious anaemia, 732

Drew, P.A. see Kiroff, G. Duckett, M. see Zaunders, J.J.

EGUCHI, K. see KURATA, A. EINHORN, S. see ROBERT, K.-H.

EISENBERG, R.A., RIBLET, R.J., LEWIS, D.E. & COHEN, P.L. Spontaneous antiidiotypic antibodies to the NZB Coomb's autoantibody, 315

ENDOH, M. see MIURA, M. ENGLISH, T.A.H. see MAHER, P. ESTES, R.G. see McLEOD, R.

FALCIONI, F., RAUTMANN, A., BERG, P.A. & GROSS, W.L. Influence of TPA (12-0-tetradodecanoylphorbol-13-acetate) on human B lymphocyte function, 613

FEIGHERY, C. see HANLY, J.G. FINN, T. see Spencer, J.

FLAD, H.-D. see HORST, H.-J.

FORD, P.M. & KOSATKA, I. Cationised IgM rheumatoid factor: in-vivo glomerular localization and immunoabsorptive capacity in the mouse, 150

FORGET, A. see BOURASSA, D. FORSBERG, P.-O. see HOLMDAHL, R.

FOULON, B. see DE WAELE, M.

FREEMAN, R.R. see PLAYFAIR, J.H. FUKUDA, T. see KURATA, A.

VAN FURTH, R. see VAN OUD ALBLAS, A.B.

GALBRAITH, R.M. see LEE, W.M.

GALLI, M. see POLI, G. GECZY, A.F. see SULLIVAN, J.S.

GEHRZ, R.C. & LEONARD, T.E. Cytomegalovirus (CMV): specific lymphokine production in congenital CMV infection, 507

GEORGOPOULOS, K., CARSON DICK, W., GOODACRE, J.A. & PAIN, R.H. A reinvestigation of the crossreactivity between Klebsiella and HLA-B27 in the aetiology of ankylosing spondylitis, 662

GERAGHTY, S.B. see VAN ASPEREN, P.P. GIORDINA, P.J. see AKBAR, A.N.

GLEESON, M. see VAN ASPEREN, P.P. GODAL, T. see MUSTAFA, A.S.

GODAL, T. see MUSTAFA, A.S. GOLDING, P.R. see GREGORY, C.D.

GOMEZ-PRIEGO, A. see PARKHOUSE, R.M.E. GOODACRE, J.A. see GEORGOPOULOS, K.

GRATAMA, J.W. see WURSCH, A.M.

GRATWOHL, A. see WURSCH, A.M. GRAY, R.W. see AKBAR, A.N.

GREGORY, C.D., LEE, H., REES, G.B., SCOTT, I.V., SHAH, L.P. & GOLDING, P.R. Natural killer cells in normal pregnancy: analysis using monoclonal antibodies and single-cell cytotoxicity assays, 121

GROSS, W.L. see FALCIONI, F. GUERAUD, V. see LARUE, C.

GULLBERG, D. see HOLMDAHL, R.

GUPTA, S. see SHARMA, S.

Haajiman, J.J. see Radl, J.

HALPERIN, M. see ROOZEMOND, R.C.

Hanly, J.G., Hassan, J., Moriarty, M., Whelan, A., Feighery, C. & Bresnihan, B. Lymphoid irradiation in intractable rheumatoid arthritis: effects on the production of immunoglobulins and rheumatoid factors, 647

HARRIS, E.N. see LOIZOU, S.

HARVEY, A.R. see PELTON, B.K. HASKARD, D.O. see YU, C.-L.

HASSAN, J. see HANLY, J.G.

HAUPTMAN, G. see MEYER, O.

HAYWARD, A.R. see TELLEZ, P.A. HELLSTROM, U.B. see SYLVAN, S.P.E.

HENRIKSSON, A., KAM-HANSEN, S. & LINK, H. IgM, IgA and IgG producing cells in cerebrospinal fluid and peripheral blood in multiple sclerosis, 176

HEYMER, B. see STRAUSS, R.

HILGARTNER, M.W. see AKBAR, A.N.

HIROFUJI, T., KAMIMOTO, K., HORI, H., NAGAI, Y., SAISHO, K., SUMIYOSHI, A. & KOGA, T. Characterization of monoclonal antibody specific for human type II collagen: possible implication in collagen-induced arthritis, 159

HODGEN, A.N. see KIROFF, G.

HOF, H. see Strauss, R. HOLDER, A.A. see Playfair, J.H.

HOLLINGSWORTH, P.N. see UKO, G.

HOLMDAHL, R., JANOSSON, L., GULLBERG, D., RUBIN, K., FORSBERG, P.-O. KLARESKOG, L. Incidence of arthritis and autoreactivity of anti-collagen antibodies after immunization of DBA/1 mice with heterologous and autologous collagen II, 639

HOLMGREN, J. see LYCKE, N.

HOLT, P.G., DEGEBRODY, A., O'LEARY, C., KRSKA, K. & PLOZZA, T. T-cell activation by antigenpresenting cells from lung tissue digests: suppression by endogenous macrophages, 586

HOLTER, W., MAIDIC, O., STOCKINGER, H. & KNAPP, W. Analysis of T cell activation with nonmitogenic anti-CD3 antibody and the phorbol ester TPA, 600

Hori, H. see Hirofuji, T.

HORST, H.-J. & FLAD, H.-D. Binding of interleukin 2 to target cells: determination of high and low affinity binding sites on T-lymphoblasts by differential dissociation, 579

Hoshino, T. see Sakane, T.

Hosking, C.S. see Pullen, G.R.

HOUGH, D. see MADDISON, P.J.

HOUTMAN, P.M., KALLENBERG, C.G.M., LIMBURG, P.C., HUITEMA, M.G., VAN RUSWIJK, M.H. & THE, T.H. Quantitation of antibodies to nucleoribonucleoprotein by ELISA: relation between antibody levels and disease activity in patients with connective tissue disease, 696

HUITEMA, M.G. see HOUTMAN, P.M.

IKARI, N. see KURATA, A.

INTRONA, M. see POLI, G.

IRVING, W.L. see AL-SAKAFF, L. ISAACSON, P.G. see Spencer, J.

ISHII, Y. see ISHIKURA, H.

ISHIKURA, H., MATSUURA, A., ISHII Y., NATORI, T.,
KIKUCHI, K. & AIZAWA, M. Different distributions of T cell subsets between perivascular and
interstitial peritubular areas during rejection of
rat renal allografts: a quantitative and ultrastructural study using monoclonal antibodies, 570

ITOH, T. see TATENO, M.

JACK, I. see McNeilage, L.J. JAMIESON, G.G. see KIROFF, G. JANOSSY, G. see PARKHOUSE, R.M.E. JANOSSON, L. see HOLMDAHL, R. JANSEN, J. see WURSCH, A.M.

JANSSENS, A.R. see VAN OUD ALBLAS, A.B.

JIRILLO, E. see ANTONACI, S.

JOHNSON, S.M. A B-cell hybridoma product inhibiting antibody secretion via 5'-nucleotidase, 412

JOHNSON, A.R. see YU, C.-L. JONES, H.M. see KNOTT, L.J.

JULIUSSON, G. see ROBERT, K.-H.

KAKIMOTO, K. see HIROFUJI, T.

KALLENBERG, C.G.M. see HOUTMAN, P.M. KAM-HANSEN, S. see HENRIKSSON, A.

KANAI, Y. see KUBOTA, T.

KANAZAWA, H. see KURATA, A. KATAMINE, S. see KURATA, A.

KAY, A.B. Eosinophils as effector cells in immunity and hypersensitivity disorders, 1

KAY, A.B. see RICHERSON, H.B.

KAY, P. see UKO, G.

KAZAMA, T., NAKAMURA, T., MORIOKA, T., OITE, T., SATO, Y. & SHIMIZU, F. A nephropathy induced by immunization of rats with EHS tumour, 104

KAZATCHKINE, M.D. see TURNER, M.W. KEMP, A.S. see VAN ASPEREN, P.P.

KIKUCHI, K. see ISHIKURA, H.

KIROFF, G. HODGEN, A.N., DREW, P.A. & JAMIESON, G.G. Lack of effect of splenic regrowth on the reduced antibody responses to pneumococcal polysaccharides in splenectomized patients, 48

KIYONO, H. see ANTONACI, S.

KLARESKOG, L. see HOLMDAHL, R. KLASTER, P.R., DE WIT, Y.L. & KOLK, A.H.J. An ELISA-inhibition test using monoclonal antibody for the serology of leprosy, 468

KNAPP, W. see HOLSTER, W.

KNOTT, L.J., LEVINSKY, R.J., NEWLAND, A.C., JONES, H.M. & LINCH, D.C. Bone marrow T cell colonyforming cells: studies of their origin and use on monitoring T cell-deplated bone marrow grafts, 561

KOGA, T. see HIROFUJI, T. KOLK, A.H.J. see KLASTER, P.R. KONDO, N. see TATENO, M. KOSATKA, I. see FORD, P.M.

KRETTLI, A.U., DE CARVALHO, L.C.P. Binding of C3 fragments to the Trypanosoma cruzi surface in the absence of specific antibodies and without prior activation of the complement cascade, 270 KRSKA, K. see HOLT, P.G.

KRZEKOTOWSKA, D. see OEN, K.

KUBOTA, T., AKATSUKA, T. & KANAI, Y. DNA affinity chromatography: application in the isolation of distinct antibody populations from SLE sera, 321

KURATA, A., KATAMINE, S., FUKUDA, T., MINE, M., IKARI, N., KANAZAWA, H., MATSUNAGA, M., EGUCHI, E. & NAGATAKI, S. Production of a monoclonal antibody to a membrane antigen of human T cell leukaemia virus (HTLV1/ATLV)-

infected cell lines from a systemic lupus erythematosus (SLE) patient: serological analyses for HTLV1 infections in SLE patients, 65

KUROKI, T., RUF, J., WHELAN, L., MILLER, A. & WALL, J.R. Antithyroglobulin monoclonal and autoantibodies cross-react with an orbital connective tissue membrane antigen: a possible mechanism for the association of ophthalmopathy with autoimmune thyroid disorders, 361

LARUE, C., GUERAUD, V. & RIVAT, C. Suitable hollow fibre immunobioreactors for specific ex vivo removal of antibodies and antigens from plasma,

LAZZARIN, A. see POLI, G.

LEE, H. see GREGORY, C.D.

LEE, W.M., MARTIN, K.L., SHELTON, L.L. & GAL-BRAITH, R.M. Hepatic membrane antibodies: studies of prevalence and specificity, 715

LEIJH, P.C.J. see VAN OUD ALBLAS, A.B.

LEONARD, T.E. see GEHRZ, R.C. LEVINSKY, R.J. see KNOTT, L.J.

LEWIS, D.E. see EISENBERG, R.A.

LIEW, F.Y. New aspects of vaccine development, 225

LIMBURG, P.C. see HOUTMAN, P.M.

LINCH, D.C. see KNOTT, L.J.

LINDHOLM, L. see LYCKE, N. LINK, H. see HENRIKSSON, A.

LOIZOU, S., McCREA, J.D., RUDGE, A.C., REYNOLDS, R., BOYLE, C.C. & HARRIS, E.N. Measurement of anti-cardiolipin antibodies by an enzyme-linked immunosorbent assay (ELISA): standardization and quantitation of results, 738

LOOF, L. see MENDEL-HARTVIG, I. LUNDBERGH, P.R. see SYLVAN, S.P.E.

LYCKE, N., LINDHOLM, L. & HOLMGREN, J. Cholera antibody production in vitro by peripheral blood lymphocytes following oral immunization of humans and mice, 39

MACKAY, I.R. see McNeilage, L.J.

MADDISON, P.J., SKINNER, R.P., VLACHOYIANNOPOU-LOS, P., BRENNARD, D.M. & HOUGH, D. Antibodies to nRNP, Sm, Ro(SSA) and La(SSB) detected by ELISA: their specificity and interrelations with rheumatoid arthritis, 346 MADDY, A.H. see Brown, V.A.

MAHER, P., O'TOOLE, C.M., WREIGHITT, T.G., SPIE-GELHALTER, D.J. & ENGLISH, T.A.H. Cytomegalovirus infection with chronic T cell subset ratio inversion with expansion of a Leu-7+ Tsc+

subset, 515 MAJDIC, O. see HOLTER, W. MANTOVANI, A. see POLI, G.

MARGOLESE, R.G. & WAINBERG, M.A. Enhanced viral inhibition of lymphocyte mitogenesis in patients with advanced breast cancer, 85

MARTIN, K.L. see LEE, W.M.

MASCART-LEMONE, F. see MEYER, O.

MASON, D.Y. see SPENCER, J. MATSUNAGA, M. see KURATA, A.

MATSUURA, A. see ISHIKURA, H.

McCrea, J.D. see Loizou, S. McGhee, J.R. see Antonaci, S.

McGrath, K. & Patterson, R. Immunology of streptokinase in human subjects, 421

McLeod, R. & Estes, R.G. Role of lymphocyte blastogenesis to *Toxoplasma gondii* antigens in containment of chronic, latent *T. gondii* infection in humans. 24

in numans, 24

McNeilage, L.J., Whittingham, S., Jack, I. & Mackay, I.R. Molecular analysis of the RNA and protein components recognized by anti-La(SS-B) autoantibodies, 685

MELLIS, C.M. see VAN ASPEREN, P.P.

MENDEL-HARTVIG, I., NELSON, B.D., LOOF, L. & TOTTERMAN, T.H. Primary biliary cirrhosis: further biochemical and immunological characterization of mitochondrial antigens, 371

MEYER, O., HAUPTMANN, G. TAPPEINER, G., OCHS, H.D. & MASCART-LEMONE, F. Genetic deficiency of C4, C2 or C1q and lupus syndromes. Association with anti-Ro (SS-A) antibodies, 678

MICHALEK, S.M. see ANTONACI, S. MIDDELDORP, J.M. see WURSCH, A.M.

MILLER, A. see KUROKI, T.

MINE, M. see KURATA, A.

MIURA, M., TOMINO, Y., YAGAME, M., ENDOH, M., SUGA, T., NOMOTO, Y. & SAKAI, H. Immunofluorescent studies of alpha-2-plasmin inhibitor (α2-PI) in glomeruli from patients with IgA nephropathy, 380

MOQBEL, R. see RICHERSON, H.B. MORIARTY, M. see HANLY, J.G. MORIOKA, T. see KAZAMA, T. MORONI, M. see POLI, G.

MOWBRAY, J.F. see TURNER, M.W.

MUSTAFA, A.S. & GODAL, T. BCG-induced suppressor T cells optimal conditions for in vitro induction and mode of action, 474

NÆSS, A., GLETTE, J., HALSTENSEN, A.I., SANDBERG, S. & SOLBERG, C.O. In vivo and in vitro effects of doxycycline on leucocyte membrane receptors, 310

310
NAESSENS, A. see DE WAELE, M.
NAGAI, Y. see HIROFUII, T.
NAGASAWA, S. see SEYA, T.
NAGATAKI, S. see KURATA, A.
NAKAMURA, T. see KAZAMA, T.
NATORI, T. see ISHIKURA, H.
NELSON, B.D. see MENDEL-HARTVIG, I.
NENCIONI, L. see TAGLIAUBE, A.
NEWLAND, A.C. see KNOTT, L.J.
NISSEN, C. see WURSCH, A.M.
NIWA, Y. see SAKANE, T.

NOMOTO, Y. see MIURA. M.
OCHS, H.D. see MEYER, O.
ODOM, L. see TELLEZ, P.A.

OEN, K., SCHROEDER, M.L. & KRZEKOTOWSKA. Pokeweed mitogen and Staphylococcus aureus Cowan I induced immunoglobulin A synthesis by lymphocytes of IgA deficient blood donors, 387

OITE, T. see KAZAMA, T.
O'LEARY, C. see HOLT, P.G.
OSTLUND, L. see ROBERT, K.-H.
O'TOOLE, C.M. see MAHER, P.
OVLAQUE, G. see SANTORO, F.

POZZILLI, P. see AL-SAKAFF, L. PAIN, R.H. see GEORGOPOULOS, K.

PARKHOUSE, R.M.E., BOFILL, M., GOMEZ-PRIEGO, A. & JANOSSY, G. Human macrophages and Tlymphocyte subsets infiltrating nodules of Onchocerca volvulus, 13

PATTERSON, R. see McGrath, K. Pedersen, J.S. see Dow, C.A. Pedini, B. see Betterle, C. Pelletier, M. see Bourassa, D.

PELTON, P.K., HARVEY, A.R. & DENMAN, A.M. The rheumatoid synovial membrane participates in systemic anti-viral immune responses, 657

PENNY, R. see ZAUNDERS, J.J.

PERI, G. see POLI, G.

PIERCE, R. see SANTORO, F.

PLAYFAIR, J.H., DE SOUZA, J.B., FREEMAN, R.R. & HOLDER, A.A. Vaccination with a purified bloodstage malaria antigen in mice; correlation of protection with T cell mediated immunity, 19

PLOZZA, T. see HOLT, P.G.

POLI, G., BOTTAZZI, B., ACERO, R., BERSANI, L., ROSSI, V., INTRONA, M., LAZZARIN, A., GALLI, M. & MANTOVANI, A. Monocyte function in intravenous drug abusers with lymphadenopathy syndrome and in patients with acquired immunodeficiency syndrome: selective impairment of chemotaxis, 136

POLI, G., INTRONA, M., ZANABONI, F., PERI, G., CARBONARI, M., AIUTI, F., LAZZARIN, A., MOR-ONI, M. & MANTOVANI, A. Natural killer cells in intravenous drug abusers with lymphadenopathy, 128

POULTER, L.W. see COLLINGS, L.A. PULFORD, K.A.F. see Spencer, J.F.

PULLEN, G.R., HOSKING, C.S. Differentiated HL60 promyelocytic leukaemia cells have a deficient myeloperoxidase/halide killing system, 304

RADL, J., VIEVEEN, M.H.M., VAN DER AKKER, T.W., BENNER, R., HAAIJMAN, J.J. & ZURCHER, C. Idiopathic paraproteinaemia V. Expression of Igh5 allotypes within the homogeneous immunoglobulins of ageing (C57BL/LiARij×CBA/ BrARij)F, mouse, 405

RAUTMANN, A. see FALCIONI, F. REES, G.B. see GREGORY, C.D. REYNOLDS, R. see LOIZOU, S. RIBLET, R.J. see EISENBERG, R.A.

RICHERSON, H.B., WALSH, G.M., WALPORT, M.J., MOQBEL, R. & KAY, A.B. Enhancement of human neutrophil complement receptors: a comparison of the rosette technique with the uptake of radio-labelled anti-CR1-monoclonal antibody, 442

VAN RUSWIJK, M.H. see HOUTMAN, P.M.

RIVAT, C. see LARUE, C.

ROBERT, K.-H., EINHORN, S., JULIUSSON, G., OST-LUND, L. & BIBERFELD, P. Interferon induces proliferation and differentiation in chronic lymphocytic leukaemia cells, 530

ROOZEMOND, R.C., HALPERIN, M. & Das, P.K. Inhibition of natural killer cell-mediated cytotoxicity by lipids extracted from Mycobacterium bovis

BCG, 482

ROSSI, V. see POLI, G. ROTHER, U. see RUPPEL, A. RUBIN, K. see HOLMDAHL, R. RUDGE, A.C. see LOIZOU, S. RUF, J. see KUROKI, T.

RUPPEL, A., DIESFELD, H.J. & ROTHER, U. Immunoblot analysis of Schistosoma mansoni antigens with sera of schistosomiasis patients: diagnostic potential of an adult schistosome polypeptide, 499

RUTTER, G. see UHLIG, H.

SAIAHO, K. see HIROFUJI, T.

SAKAI, H. see MIURA, M.

SAKANE, T., UEDA, Y., SUZUKI, N., NIWA, Y., HOS-HINO, T. & TSUNEMATSU, T. OKT4\* and OKT8\* T lymphocytes produce soluble factors that can modulate growth and differentiation of human B cells, 112

SALMON, M., BACON, P.A., SYMMONS, D.P.M. & BLANN, A.D. Transferrin receptor bearing cells in the peripheral blood of patients with rheumatoid arthritis. 346

SALVIATI, C. see BETTERLE, C.

SANTORO, F., AFCHAIN, D., PIERCE, R., CESBRON, J.Y., OVLAQUE, G. & CAPRON, A. Serodiagnosis of toxoplasma infection using a purified parasite protein (P30), 262

SATO, Y. see KAZAMA, T. SAVA, P. see VUITTON, D.A.

SCHROEDER, M.L. see OEN, K. SCOTT, I.V. see GREGORY, C.D.

SEILLES, E. see VUITTON, D.A. SENSI, M. see AL-SAKAFF, L.

SEYA, T., NAGASAWA, S. & ATKINSON, J.P. Generation of the bioreactive kallikrein-derived fragment, C3d-k, by HANE plasma, 208

SEYMOUR, N.D. see TURNER, M.W.

SHAH, L.P. see GREGORY, C.D.

SHARMA, B. & GUPTA, S. Antigen-specific primary cytotoxic T lymphocytes (CTL) responses in acquired immunodeficiency syndrome (AIDS) and AIDS-related complexes (ARC), 296

SHELTON, L.L. see LEE, W.M. SHIMIZU, F. see KAZAMA, T.

Shinomiya, N. & Yata, J. In-vitro analysis of defective IgA production in selective IgA deficiency in childhood, 746

SKAMENE, E. see BOURASSA, D.

SKINNER, R.P. see MADDISON, P.J. SLOANE, J.P. see DILLY, S.A.

SMITH, S.K. see Brown, V.A. SMITH, J.B. see BOCCHIERI, M.H.

SMITH JR, J.B. see BOCCHIERI, M.H.

SPECK, B. see WURSCH, A.M.
SPENCER, J., FINN, T., PULFORD, K.A.F., MASON, D.Y. & ISAACSON, P.G. The human gut contains a novel population of B lymphocytes which resemble marginal zone cells, 607

SPIEGELHALTER, D.J. see MAHER, P. STARUCH, M.I. see BOCCHIERI, M.H.

STOCKDILL, G. see Brown, V.A. STOCKINGER, H. see HOLTER, W.

STOHL, W. Modulation of the immune response by immunoglobulin for intravenous use. I. Inhibition of pokeweed mitogen-induced B cell differentiation, 220

STRAUSS, R., HEYMER, B. & Hof, H. Effects of cyclosporin A on experimental infection with Listeria monocytogenes, 491

SUGA, T. see MIURA, M.

SULLIVAN, J.S. & GECZY, A.F. The modification of HLA-B27-positive lymphocytes by the culture filtrate of Klebsiella K43 BTS 1 is a metabolically active process, 672

SUMIYOSHI, A. see KOGA, T.

SUZUKI, N. see SAKANE, T.

SYLVAN, S.P.E., HELLSTROM, U.B. & LUNDBERGH, P.R. Detection of cellular and humoral immunity to hepatitis B surface antigen (HBsAg) in asymptomatic HBsAg carriers, 288

SYMMONS, D.P.M. see SALMON, M.

SYPEK, J.P. & WYLER, D.J. Cell contact-mediated macrophage activation for anti-leishmanial defence: mapping of the genetic restriction to the I region of the MHC, 449

TAGLIAUBE, A., NENCIONI, L., CAFFARENA, A., VILLA, L., BORASCHI, D., CAZZOLA, G. & GALVALIERI, S. Cellular immunity against Salmonella typhi after live oral vaccine, 242

TAPPEINER, G. see MEYER O.

TATENO, M., KONDO, N. & YOSHIKI, T. Autoimmune disease and malignant lymphoma associated with host-versus-graft disease in mice, 535

TELLEZ, P.A., ODOM, L. & HAYWARD, A.R. Immunity to herpes simplex virus in children receiving treatment for acute lymphoblastic leukaemia (ALL), 525

THE, T.H. see WURSCH, A.M.

---, see HOUTMAN, P.M.

TOH, B.H. see Dow, C.A. TOMINO, Y. see MIURA, M.

TOTTERMAN, T.H. see MENDEL-HARTVIG, I.

TRON, F. see CUKIER, R.

TSUNEMATSU, T. see SAKANE, T.

TURCOTTE see BOURASSA, D.

TURNER, M.W., SEYMOUR, N.D., KAZATCHKINE, M.D. & MOWBRAY, J.F. Suboptimal C3b/C3bi deposition and defective yeast opsonization. I. Evidence for the absence of essential co-factor activity, 427

TURNER, M.W., SEYMOUR, N.D., KAZATCHKINE, M.D. & MOWBRAY, J.F. Suboptimal C3b/C3bi deposition and defective yeast opsonization. II. Partial purification and preliminary characterization of an opsonic co-factor able to correct sera with the defects, 435

UEDA, Y. see SAKANE, T.

UHLIG, H., RUTTER, G. & DERNICK, R. Self-reactive B lymphocytes detected in young adults, children and newborns after in vitro infection with Epstein-Barr virus, 75

UKO, G., DAWKINS, R.L., KAY, P., CHRISTIANSEN, F.T. & HOLLINGSWORTH, P.N. CR1 deficiency in SLE: acquired or genetic? 329

UNGAR, B. see Dow, C.A.

VIEVEEN, H.M. see RADL, J.

VILLA, L. see TAGLIAUBE, A.

VLACHOYIANNOPOULOS, P. see MADDISON, P.J.
VUITTON, D.A., SEILLES, E., CLAUDE, P., SAVA, P. &
DELACROIX, D. I. Gall bladder: the predominant

DELACROIX, D.L. Gall bladder: the predominant source of bile IgA man, 185

- WAINBERG, M.A. see MARGOLESE, R.G.
- WALL, J.R. see KUROKI, T.
- WALPORT, M.J. see RICHERSON, H.B.
- WALSH, G.M. see RICHERSON, H.B.
- WATERS, M.F.R. see Collings, L.A.
- WEILAND, E. & WEILAND, F. Detection and preliminary characterization of a transmissible agent inducing autoantibody against Golgi-antigen, 167
- WEILAND, F. see WEILAND, E.
- WHELAN, L. see KUROKI, T. WHELAN A. see HANLY, J.G.
- WHITTINGHAM, L.J. see MCNEILAGE, L.J.
- WILLIAMSON, S.I. see ANTONACI, S.
- WOOD, D.D. see BOCCHIERI, M.H.
- WREIGHT, T.G. see MAHER, P.
  WURSCH, A.M., GRATAMA, J.W., MIDDELDORP, J.M.,
  NISSEN, C., GRATWOHL, A., SPECK, B., JANSEN, J.,
  D'AMARO, J.D., THE, T.H. & DE GAST, G.C. The
  effect of cytomegalovirus infection of T lympho-

cytes after bone marrow transplantation, 278

- WYLER, D.J. see SYPEK, J.P.
- YAGAME, M. see MIURA, M.
- YATA, J. see SHINOMIYA, N.
- YOSHIKI, T. see TATENO, M.
- Young, Y. see Zaunders, J.J.
- YU, C.-L., HASKARD, D.O., CAVENDER, D., JOHNSON, A.R. & ZIFF, M. Human gamma interferon increases the binding of T lymphocytes to endothelial cells, 554
- ZANABONI, F. see POLI, G.
- ZANABONI, F. See POLI, G.
  ZAUNDERS, J.J., COOPER, D.A., YOUNG, Y., DUCKETT,
  M., PENNY, R. & ZIEGLER, J.B. Suppression of IgE synthesis in vitro by allogeneic T cells from atopic and non-atopic subjects, 193
- ZEVIANI, M. see BETTERLE, C.
- ZIEGLER, J.B. see ZAUNDERS, J.J.
- ZIFF, M. see Yu, C.-L.
- ZURCHER, C. see RADL, J.

## SUBJECT INDEX

Acquired immunodeficiency syndrome, 128, 136, 296 Complement Affinity column chromatography (DNA), 321 Age-linkage, 632 Ageing, 405 Alcoholic cirrhosis, 724 Allergy, 1 Androgen, 630 Ankylosing spondylitis, 662, 672 Antibodies anti-CD3, 600 anti-DNA, 321, 329 anti-La(SSB), 337 anti-nRNP, 337 anti-phospholipid, 738 anti-Ro, 337, 678 anti-Sm, 337 suppression, 412 Anticardiolipin, 738 Antigen presentation, 586 Antigens FX1A-related, 104 enteric, 672 mitochondrial, 371 schistosome Antiidiotypic antibodies, 315 Arthritis (collagen induced) 159, 639 Asymptomatic HBsAg carriers, 288 Atopic disease, 753 Atopic state, 193 Atopy, 753 Autoantibodies, 75, 167 anti-La(SSB), 685 anti-parathyroid, 353 Coombs, 315 mitochondrial, 353 Autoimmunity, 535, 622, 630

B cell
differentiation, 112, 200
factor, 412
growth factor, 112, 748
B lymphocytes, 75, 607
BCG, 474
Benign monoclonal gammapathy, 405
Bile, 185
Bone marrow T cell colony forming cells, 377
Bone marrow transplantation, 278, 545
Breast cancer, 85

C3 fragments, 270, 427, 435
C3b receptor, 310
C3b/C3bi, 427, 435
C3d generation, 208
C3d,g, 208
C3d,g, 208
Cell mediated immunity, 19, 31, 630
Cellular immunity, 242
Cerebrospinal fluid, 176
Chemotaxis, 136
factor (fMLP), 442
Cholera, 39
Chrysotherapy, 705
Collagen type II, 159, 639

Complement activation, 270, 329 deficiency, 672 receptor deficiency, 329 receptor enhancement, 442 Connective tissue disease, 337, 696 Cyclosporin A, 491 Cytoadherence, 248 Cytomegalovirus, 278, 507, 515 Cytotoxicity assay, 121

DNA, 321 Delayed type hypersensitivity, 31, 491 Dendritic cells, 458 Differentiation, 304, 530 Doxycycline, 310

Endothelial cells, 554
Enzyme linked immunosorbent assay, 321, 337, 468, 662, 696, 738
Eosinophils, 1
Epstein-Barr virus, 75

F<sub>1</sub>-ATPase, 371 F<sub>1</sub> hybrids, 315 Familial disease, 57 Familial hypercholesterolemia, 217 Fc receptor, 310, 412 Fibrinogen, 380

Gall bladder bile, 185
Gammaglobulin, 200
Gastritis, 732
Gene Beg, 31
Genetic deficiency, 678
Genetic restriction, 449
Glomerulus, 150
Golgi antibody, 167
Graft versus host disease, 535, 545, 561
Gram-negative bacteria, 248
Granulocytes, 57, 724
Granuloma, 491
Graves' disease, 361
Graves' disease, 361
Graves' ophthalmopathy, 361
Gut, 607

Helminths, I
Hepatic membrane antibodies, 715
Hepatitis B surface antigen, 288
Hereditary angioneurotic edema, 208
Herpes simplex virus, 525
Heymann nephritis, 104
Hodgkin's disease, 24
Host defence, 449
Human macrophages, 13
Hypercalcaemia, 57
Hypersensitivity, I

Idiopathic hypoparathyroidism, 353 paraproteinaemia, 405 Idiotype, 225 IgA bile, 185 deficiency, 387, 748, 753 nephropathy, 380 synthesis, 387 IgE synthesis, 193 IgG serodiagnosis, 262 IgM serodiagnosis, 262 Immune complexes cationic, 150 Immune response, 225 Immunelectron microscopy, 570 Immunity 1, 31, 491, 525 Immunization, 104 Immunoabsorption, 150 Immunoadsorption therapy, 217 Immunobioreactors hollow fibre, 217 Immunoblot analysis, 499, 732 Immunodiagnosis, 499 Immunofluorescence, 380 Immunoglobulin producing cells, 176 Immunoglobulin synthesis, 248 Immunoglobulins, serum/cytoplasmic, 397 Immunohistology, 545 Immunological characterization, 371 Immunosuppression, 13, 515 Infiltration, 13 Interferon, 530, 554 Interleukin 1, 474, 622, 705 Interleukin 2, 112, 296, 474, 622, 705 receptor, 579 Interstitial lung disease, 49 Intravenous drug abusers, 128, 136

Kallikrein, 208 Klebsiella, 662, 672

Lectins, 248 Leishmaniasis, 449 Leprosy lepromatous, 458 tuberculoid, 458 Leprosy serology, 468 Leu-7+ Ts-c+ subset, 515 Leucocyte, 310 polymorphonuclear, 304 Leukaemia B chronic lymphocytic, 95 adult T cell, 65 lymphoblastic, 525 lymphocytic, 530 promyelocytic, 304 human T cell, 65 Leukaemic phenotype, 95 Listeria monocytogenes, 491 Liver, 545, 715 Lung, 586 Lupus nephritis- DNA, 143 Lupus syndromes, 678 Lymphadenopathy, 256, 535 Lymphadenopathy syndrome, 128, 136 Lymphocyte blastogenesis, 24 mitogenesis, 85 subpopulations, 545 Lymphocytes activated, 346

Natural killer, 594 peripheral blood, 39 Lymphoid irradiation, 647 Lymphokines, 507 Lymphoma, 535 Macrophages, 13, 449, 586 Major Histocompatibility Complex, 449 Malaria, 19 Marginal zone cells, 607 Membrane antigen, 65 fluidity, 482 receptors, 310 CBA, 315 NZB, 315 NZB × C58 recombinant, 622 Mitochondrial antigens, 353, 371 autoantibodies, 353 Mitogen activity, 613 Mixed lymphocyte culture, 594 Monoclonal antibodies, 65, 95, 143, 159, 256, 278, 361, 442, 468, 570, 594 Monocytes, 724 Multiple sclerosis, 176 Mycobacterial lipids, 482 Mycobacterium bovis, 31 Mycobacterium leprae, 458 Myeloperoxidase, 304 Nandrolone decanoate, 630 Natural killer cells, 121, 128, 482

Neutrophil, 442 function, 304 5'-nucleotidase, 412

Onchocerciasisi, 13 Opsonic cofactor, 427, 435 Opsonization, 427, 435 Oral immunization, 39 Oral vaccine, 242

Parathyroid hormone, 57 Pemphigus vulgaris, 705 Peptides, 225 Peripheral blood lymphocytes, 39 Peripheral blood mononuclear cells, 193 Pernicious anaemia, 732 Peyer's patches, 607 Phytohaemaglutinin, 85, 594 Plaque-forming cells, 200 Plasma cell, 412 a2-plasmin inhibitor, 380 Pneumococcal polysaccharides, 48 Pokeweed mitogen, 200, 387, 594 Pregnancy (normal), 121 Primary biliary cirrhosis, 321, 724 Protein (P30), 262

RNA, 685 SnRNP, 685 Renal transplantation, 570 Rheumatoid arthritis, 346, 647, 657 Rheumatoid factor, 150, 647 Rheumatoid synovial membrane, 657

## Subject Index

Salmonella minnesota, 248 Salmonella typhi, 242 Schistomiasis, 499 Schistosoma mansoni, 499 Secretory component, 185 Self tolerance, 75 Septicaemia, 49 Sex-linkage, 639 Sjögrens syndrome, 685 Splenectomy, 49 Splenosis, 48 Staphylococcus aureus, 387 Streptokinase, 421 Surface glycoproteins, 95 Synovectomy, 657 Systemic lupus erythematosus, 65, 143, 321, 329, 337, 535

T cell activation, 288, 586, 600 colonies, 561 growth factor, 85 helper factor, 748 leukaemia viruses (human), 65 subsets, 112, 248, 570 T cells allogeneic, 193 suppressor, 474

T lymphoblasts, 579 T lymphocytes, 278, 254 subsets, 13 T-cytotoxic/suppressor cells, 256 TPA(12-0-tetradecanylphorbol-13-acetate), 600, 613 Thalassaemia, 397 Thyroglobulin, 361 Thyroid, 361 Toxoplasma gondii, 24, 256, 262 Toxoplasmosis, 256, 262 Transferrin receptors, 346 Transfusion, 397 Transmissible agent, 167 Trypanosoma cruzi, 270 Trypomastigotes, 270 Tuberculosis, 474 Tumour (EHS), 104 Typhoid fever, 242

Typhoid fever, 242

Vaccine, 19
development, 225
influenza virus, 657
oral, 242

Vaccine, polyvalent pneumococcal polysaccharide, 48

Viral inhibition, 85

Yeast, 427, 435

